# *Global Tsunami Impacts: A Comprehensive Study of Death and Damage Patterns*

Yourfee Kennedy, Biosphere Reserves Management. Eberswalde University for Sustainable Development.

## Abstract

This exhaustive investigation explores the intricate patterns and dynamics of tsunami effects on a global scale, with a specific focus on the overall number of fatalities and the extent of damages. Tsunamis, which are triggered by seismic activities, are renowned for their catastrophic nature and their ability to cause extensive devastation beyond the confines of coastal areas. The research draws insights from a comprehensive review of the existing literature, shedding light on the multifaceted ramifications of these natural disasters and underscoring the importance of a nuanced understanding. The methodology entails the acquisition of reliable and comprehensive data from reputable databases, such as NOAA's National Centers for Environmental Information and the Pacific Tsunami Warning Canter. To analyze the spatial dimensions of tsunami impacts, map the total number of casualties and the extent of damages, and identify distribution patterns, Geographic Information System (GIS) tools are employed. The study conducts thorough examinations of specific instances of tsunamis to provide a contextual understanding of the factors that influence the severity of the consequences. The results, which are illustrated in Figures 1 to 5, offer valuable insights into the consequences of different tsunami events, including their impact, total damage, and death tolls. The discussion section provides an interpretation of these results, highlighting the diverse nature of tsunamis and the varied outcomes they produce in different regions. The analysis underscores the importance of tailored interventions and preparedness measures that are specific to each region, taking into account the severity of the tsunami. In conclusion, this study places great emphasis on the critical significance of early warning systems, resilient infrastructure, and community preparedness in mitigating the effects of tsunamis. The findings contribute to a broader comprehension of these natural disasters, serving as a basis for future research and the formulation of effective policies. Given the paramount importance of continuous efforts in research, technology, and community engagement, this study serves as a valuable resource for enhancing global resilience to the evolving nature of tsunamis.

## 1 Introduction

Tsunamis, characterized by immense ocean waves triggered by seismic activities, have been proven to be catastrophic natural phenomena with wide-ranging consequences. The devastation caused by tsunamis extends beyond immediate coastal areas, affecting human lives and resulting in significant economic losses. The comprehension of the patterns and dynamics of tsunami impacts is vital for efficient disaster management and mitigation. This research embarks on a thorough exploration of the global effects of tsunamis, with a specific focus on the crucial factors of total fatalities and damages. By delving into the intricacies of these impacts, the study aims to provide valuable insights that can guide proactive measures to enhance global resilience to tsunami disasters (Bolt 2004; de Boer and Sanders 2005; Guidoboni & Ebel,2009; Plag, 2014; Scourse et al. 2018).

## 2 Literature Review

The literature regarding the impacts of global tsunamis accentuates the necessity for a nuanced comprehension of the multifaceted ramifications associated with these natural calamities. Investigations conducted by various researchers (Tetsuka et al. 2020) underscore the significance of evaluating the aggregate loss of life and destruction as primary indicators of the severity of tsunami incidents, thus providing essential insights into the magnitude of these disasters. The examination of temporal and spatial patterns of tsunami impacts on a global scale has been undertaken by (Harbitzet al. 2013), elucidating the variability in the frequency and intensity of these events. Through the analysis of historical data, these studies contribute to a comprehensive comprehension of the factors that influence the total loss of life and destruction, as well as identifying regions and periods that are particularly vulnerable. The investigation of the human and economic consequences of tsunamis has been a central focus for researchers (Pagnoni et al.,2020). By exploring the correlation between the total loss of life and destruction, these studies uncover the potential socioeconomic impacts of these natural disasters. interconnected nature of these impacts, with socio-economic factors often exacerbating vulnerabilities and influencing the magnitude of disasters (Tinti et al., 2011; Suppasri etal., 201).

## 3 Methodology

The foundation of this study is rooted in the acquisition of reliable and comprehensive data regarding the occurrences of tsunamis worldwide. To gather historical records of tsunamis, which encompass information on the total number of casualties and the economic consequences, credible databases such as NOAA's National Centers for Environmental Information (NCEI) and the Pacific Tsunami Warning Center will be consulted. The selected dataset will cover a significant timeframe to encompass a diverse range of events. To examine the spatial dimensions of tsunami impacts, Geographic Information System (GIS) tools will be employed. The data concerning the total number of casualties and damages will be visually represented through mapping, which will facilitate the identification of distribution patterns across various geographical regions. Additionally, spatial clustering and hotspots will be determined, providing valuable insights into the susceptibilities of different areas to tsunamis. To enrich the analysis, in-depth examinations of specific tsunami occurrences will be conducted. These specific examinations will provide a contextual understanding of the elements that affect the magnitude of consequences, such as local geographical features, early warning systems, and preparedness measures (Fraser et al. 2009, 2010).

The research will conclude with the formulation of recommendations for focused mitigation strategies based on the observed trends. By identifying areas that are susceptible and comprehending the intricate interplay of factors that contribute to overall casualties and destruction, this investigation seeks to enlighten policymakers and authorities in the field of disaster management regarding the creation of effective measures to bolster global resilience to the impacts of tsunamis (Davies et al. 2017).

## 4 Results


Global Tsunami Impact: Total Damage and Categorization
(Tsunami 10) - Graduated Equal Count Interval
0°00′
90°00′E
180°00′
90°00′N
Moderate (~$1 to $5 million)
Severe (~$5 to $25 million)
Moderate (~$1 to $5 million)
Severe (~$5 to $25 million)
Moderate (~$1 to $5 million)
Very Many (~1,001 or more)
Very Many (~1,001 or more)
Severe (~$5 to $25 million)
Severe (~$5 to $25 million)
Moderate (~$1 to $5 million)
Very Many (~1,001 or more)
Severe (~$5 to $25 million)
Moderate (~$1 to $5 million)
Limited (roughly corresponding to less than $1 million)
Severe (~$5 to $25 million)
Limited (roughly corresponding to less than $1 million)
Severe (~$5 to $25 million)
1:52,607,366
Legend
tsunami_10
3,2000 - 5,8000
5,8000 - 6,2000
6,2000 - 6,5000
6,5000 - 6,6000
6,6000 - 6,8000
6,8000 - 6,9000
6,9000 - 7,0000
7,0000 - 7,1000
7,1000 - 7,2000
7,2000 - 7,4000
7,4000 - 7,5000
7,5000 - 7,7000
7,7000 - 7,9000
7,9000 - 8,2000
8,2000 - 9,5000

**Figure 1 shows the impact and total damage of Tsunami 10**



**Figure 2 shows the impact and total damage of Tsunami 13 and total death in thousand**


Global Tsunami Impact: Total Damage and Categorization
(Tsunami 14) - Graduated Equal Count Interval
0°00′
90°00′E
180°00′
90°00′N
Limited (roughly corresponding to less than $1 million)
Moderate (~$1 to $5 million)
Moderate (~$1 to $5 million)
Limited (roughly corresponding to less than $1 million)
Moderate (~$1 to $5 million)
Severe (~$5 to $25 million)
Very Many (~1,001 or more)
Severe (~$5 to $25 million)
Moderate (~$1 to $5 million)
Severe (~$5 to $25 million)
Moderate (~$1 to $5 million)
Moderate (~$1 to $5 million)
Moderate (~$1 to $5 million)
Severe (~$5 to $25 million)
Moderate (~$1 to $5 million)
Legend
tsunami _14
● 5,0000 - 13,1900
● 13,1900 - 18,5000
● 18,5000 - 31,5960
● 31,5960 - 34,8084
● 34,8084 - 37,0500
● 37,0500 - 38,4000
● 38,4000 - 40,5000
● 40,5000 - 43,5000
● 43,5000 - 51,4440
● 51,4440 - 71,8130
1:36,572,464

**Figure 3 shows the impact and total damage of Tsunami 14**


Global Tsunami Impact: Total Damage and Categorization
(Tsunami 15) - Graduated Equal Count Interval
0°00′
90°00′E
180°00′
90°00′N
Moderate (~$1 to $5 million)
Moderate (~$1 to $5 million)
Severe (~$5 to $25 million)
Severe (~$5 to $25 million)
Moderate (~$1 to $5 million)
Severe (~$5 to $25 million)
Moderate (~$1 to $5 million)
Limited (roughly corresponding to less than $1 million)
Limited (roughly corresponding to less than $1 million)
Limited (roughly corresponding to less than $1 million)
1:52,607,366
Legend
tsunami_15
2,2320 - 16,1600
16,1600 - 26,2220
26,2220 - 94,5533
94,5533 - 120,5000
120,5000 - 126,5000
126,5000 - 135,2000
135,2000 - 141,5000
141,5000 - 147,0000
147,0000 - 162,0163
162,0163 - 179,9840

**Figure 4 shows the impact and total damage of Tsunami 15**


Global Tsunami Impact: Total Damage and Categorization
(Tsunami 20) - Graduated Equal Count Interval
0°00′
90°00′E
180°00′
Limited (roughly corresponding to less than $1 million)
Severe (~$5 to $25 million)
Very Many (~1,001 or more)
Moderate (~$1 to $5 million)
Very Many (~1,001 or more)
Very Many (~1,001 or more)
90°00′N
90°00′N
Moderate (~$1 to $5 million)
Severe (~$5 to $25 million)
Limited (roughly corresponding to less than $1 million)
Moderate (~$1 to $5 million)
0°00′
0°00′
90°00′E
1:50,430,509
180°00′
0°00′
Legend
tsunami_20
● 1,0000 - 1,0000
● 1,0000 - 1,0000
● 1,0000 - 1,5000
● 1,5000 - 2,0000
● 2,0000 - 2,0000
● 2,0000 - 3,0000
● 3,0000 - 3,0000
● 3,0000 - 3,0000
● 3,0000 - 4,0000
● 4,0000 - 9,0000

**Figure 5 shows the impact and total damage of Tsunami 20**

## 5 Discussion

Figure 1 provides insights into the impact and total damage caused by Tsunami 10. The recorded impact ranges from 8.2 to 9.5(82,000 to 95,000 death tolls), indicating the severity of the tsunami in affected areas. Correspondingly, the damage is categorized into severe (5 million dollars or more), very many (between 1 and 5 million dollars), and limited (roughly less than 1 million dollars). The variability in impact and damage underscores the heterogeneous nature of tsunamis and their diverse consequences in different regions. In Figure 2, the maximum death toll attributed to Tsunami 13 is 4 thousand, while the minimum is 1 thousand. The locations affected, including Cutro, Rockaway Beach, NY, Cap-Haitain, and Mount Pelee, among others, experienced varying degrees of devastation. The data highlights the critical need for preparedness and early warning systems in vulnerable coastal regions to mitigate the loss of life during tsunami events. Tsunami 14, depicted in Figure 3, showcases a significant range in impact, from 5 to 71.81(maximum death toll 718,000). The corresponding damage categories include severe (5 million to 25 million dollars), very many (1 to 5 million dollars), moderate (1 to 5 million dollars), and limited (roughly less than 1 million dollars). This variation suggests the complex dynamics of tsunamis, with some areas experiencing more substantial impacts than others. Figure 4 illustrates the corresponding damage categories. Understanding the spatial distribution of impact and damage is crucial for emergency response and recovery efforts, as different regions may require tailored interventions based on the severity of the tsunami. Lastly, Figure 5 outlines the impact variation of another tsunami event, ranging from (40,000 to 90,000 death toll). The associated damage includes severe, moderate, very many, and limited. This diversity in impact and damage underscores the intricate interplay of factors influencing the consequences of tsunamis (Kawamura et al. 2020, Dilek 2017, Reese et al., 2007).

## 6 Conclusion

In conclusion, the comprehensive analysis of multiple tsunami events depicted in Figures 1 to 5 underscores the need for an in-depth understanding of these natural disasters' complex dynamics and consequences. The spatial distribution of impact, total damage, and death tolls across different events showcases tsunamis' diverse nature, emphasizing the need for region-specific preparedness and response strategies. The findings from this study reinforce the critical importance of early warning systems, especially in coastal areas prone to tsunamis. Tsunami 10, illustrated in Figure 1, exemplifies the variability in impact and damage, highlighting the significance of real-time data and accurate predictions to mitigate the potential devastation. Figure 2 delves into the specific case of Tsunami 13, revealing variations in death tolls across different locations. This emphasizes the need for tailored evacuation plans and emergency response measures, considering the unique geographical and demographic characteristics of each affected region. Tsunami 14, as depicted in Figure 3, further emphasizes the heterogeneity in impact and damage. The wide-ranging consequences, from severe to limited, underscore the challenges faced by emergency responders in addressing the aftermath of such events. These insights should inform the development of robust disaster management strategies that account for the diverse impacts of tsunamis on different communities. Figure 4, focusing on Tsunami 15, reinforces the spatial disparities in impact and damage. The data supports the argument for investment in resilient infrastructure and community preparedness programs, acknowledging the varying degrees of vulnerability across different coastal regions. Lastly, Figure 5 provides a glimpse into yet another tsunami event, showcasing the diverse impact categories and associated damages. This reaffirms the need for ongoing research, monitoring, and adaptive strategies to address the evolving nature of tsunamis and their potential impacts on coastal communities. As we navigate the complexities of natural disasters, continuous efforts in research, technology, and community engagement are paramount. The findings of this study contribute to the broader understanding of tsunamis and their consequences, serving as a foundation for future research and the development of effective policies (Bai, Y et al., 2016, Cabral 2020, Tinti et al., 2011; Suppasri et al., 2013).

## 7 References

Bai, Y.; Adriano, B.; Mas, E.; ; Ko Shimura, S. Identifying Building Damage Patterns in the 2016 Meinong, Taiwan Earthquake Using Post-Event Dual-Polarimetric ALOS-2/PALSAR-2 Imagery. J. Disaster Res. 2018,13, 291–302.

Bai, Y.; Adriano, B.; Mas, E.; Ko Shimura, S. Machine Learning Based Building Damage Mapping from the ALOS-2/PALSAR-2 SAR Imagery: Case Study of 2016 Kumamoto Earthquake. J. Disaster Res. 2017,12, 646–65.

Bovolo, F.; Bruzzone, L. A Split-Based Approach to Unsupervised Change Detection in Large-Size Multitemporal Images: Application to Tsunami-Damage Assessment. IEEE Trans. Geosci. Remote Sens .2007, 45, 1658–1670.

Chen, S.; Sato, M. Tsunami Damage Investigation of Built-up Areas Using Multitemporal Spaceborne Full Polarimetric SAR Images. IEEE Trans. Geosci. Remote Sens. 2013, 51, 1985–1997.

Fernandez Galarreta, J.; Kerle, N.; Gerke, M. UAV-based urban structural damage assessment using object-based image analysis and semantic reasoning. Nat. Hazards Earth Syst. Sci. 2015, 15, 1087–1101.

Gupta, H. K. and Gahalaut, V. K.: Three Great Tsunamis: Lisbon (1755), Sumatra-Andaman (2004) and Japan (2011), Springer Dordrecht Heidelberg New York London, doi:10.1007/978-94-007-6576-4, 2013.

Gokon, H.; Ko Shimura, S.; Megur, K. Verification of a method for estimating building damage in extensive tsunami-affected areas using L-band SAR data. J. Disaster Res. 2017, 12, 251–258.

Hayashi, Y. Extracting the 2004 Indian Ocean tsunami signals from sea surface height data observed by satellite altimetry. J. Geophysics. Res. 2007, 113, C01001.

Hoshi, T.; Murao, O.; Yoshino, K.; Yamazaki, F.; Estrada, M. Post-Disaster Urban Recovery Monitoring in Pisco After the 2007 Peru Earthquake Using Satellite Image. J. Disaster Res. 2014, 9, 1059–1068.

Ko Shimura, S.; Hayashi, S. Tsunami flow measurement using the video recorded during the 2011 Tohoku tsunami attack. In Proceedings of the 2012 IEEE International Geoscience and Remote Sensing Symposium, Munich, Germany, 22–27 July 2012; pp. 6693–66.

Liu, W.; Yamazaki, F.; Gokon, H.; Koshimura, S. Extraction of Tsunami-Flooded Areas and Damaged Buildings in the 2011 Tohoku-Oki Earthquake from TerraSAR-X Intensity Images. Earthq. Spectra 2013, 29, S183–S200.

Liu, W.; Yamazaki, F.; Gokon, H.; Koshimura, S. Extraction of Damaged Buildings due to the 2011 Tohoku, Japan Earthquake Tsunami. In Proceedings of the 2012 IEEE International Geoscience and Remote Sensing Symposium, Munich, Germany, 22–27 July 2012; pp. 4038–4041.

Marchant, M. Advanced Remote Sensing Technology for Tsunami Modelling and Forecasting; CRC Press: Boca Raton, FL, USA, 2018; 316p.

Peek-Asa, C., Ramirez, M., Seligson, H., & Shoaf, K. (2003). Seismic, structural, and individual factors associated with earthquake-related injury. Injury Prevention: Journal of the International Society for Child and Adolescent Injury Prevention, 9(1), 62-66.

Sato, M.; Chen, S.-W.; Satake, M. Polarimetric SAR Analysis of Tsunami Damage Following the March 11, 2011 East Japan Earthquake. Proc. IEEE 2012, 100, 2861–2875.

Sullivan, K. M., & Hossain, S. M. (2010). Earthquake mortality in Pakistan. Disasters, 34(1), 176-183.

US Geological Survey. 2009. “Historic Earthquakes and Earthquake Statistics: Where do earthquakes occur?” Retrieved 12/7/09 at http://earthquake.usgs.gov/learn/faq/?categoryID=11&faqID=95

Vu, T.T.; Matsuoka, M.; Yamazaki, F. Dual-scale approach for detection of tsunami-affected areas using optical satellite images. Int. J. Remote Sens. 2007, 28, 2995–3011.

Wen, J., Shi, Y. K., Li, Y. P., Wang, L., Cheng, L., Gao, Z., et al. (2009). Risk factors of earthquake inpatient death: A case-control study. Critical Care, 13.

White, R. A., & Harlow, D. H. (1993). Destructive upper-crustal earthquakes in America since 1900. Bulletin of the Seismological Society of America, 83(4), 1115-1142.

Wisner B, Blaikie P, Cannon T, Davis I.(2008). At Risk: Natural hazards, people’s vulnerability, and disasters (2nd Edition). New York: Routledge, 275.

Yamazaki, F.; Matsuoka, M. Remote Sensing Technologies in Post-disaster Damage Assessment. J. Earthquake Tsunami 2007, 1, 193–210.